

Minimal forms for function fields
of Pfister forms

D.W. HOFFMANN

MINIMAL FORMS FOR FUNCTION FIELDS OF PFISTER FORMS

Detlev W. Hoffmann

Abstract. Let F be a field of characteristic $\neq 2$, $\{\rho_i\}$ a set of Pfister forms over F and $K = F(\{\rho_i\})$ the iterated function field of the ρ_i 's over F . We investigate the isotropy behaviour of quadratic forms over F under this field extension. Elman-Lam-Wadsworth have shown that if the Hasse number \tilde{u} of F is $\leq 2^n$, $n = 1, 2$, and if all the ρ_i are of fold $\geq n$, then K/F is excellent and the Witt kernel $W(K/F)$ is generated by $\{\rho_i\}$. We extend and generalize these results. We show, for example, that if $\tilde{u}(F) \leq 6$ or if F is linked and all the ρ_i 's are of fold ≥ 3 , then K/F is excellent. More generally, if $\tilde{u}(F) \leq 2^n$ and all the ρ_i 's are of fold $\geq n + 1$ (resp. $\geq n$), then K/F is excellent (resp. $W(K/F)$ is generated by $\{\rho_i\}$). We also investigate so-called K -minimal forms, i.e. anisotropic forms over F which become isotropic over K but no proper subform does. In two examples, one over $\mathbf{R}(t)$ and one over a formally real global field, we use the methods developed in this paper to compute the K -minimal forms explicitly.

1 Introduction

Let F be a field of characteristic $\neq 2$. In this paper we want to investigate the behaviour of quadratic forms over F over extensions K/F where K is a function field of the form $F(\{\rho_i\})$ of Pfister forms $\{\rho_i\}$. We will call such a field extension a *Pfister extension*. Extensions of this type have been given a thorough treatment by Elman, Lam and Wadsworth in [ELW 1], [ELW 2] and [ELW 3]. They focussed on the following questions: What is the Witt kernel $W(K/F)$ of the ring homomorphism $WF \rightarrow WK$ between the Witt rings of F and K induced by scalar extension? Is the extension K/F excellent, i.e., does there exist to any form φ over F a form ψ over F such that the anisotropic part of φ over K , $(\varphi_K)_{\text{an}}$, is isometric to ψ_K ?

Our main motivation will be the determination of so-called K -minimal forms for such extensions. These are anisotropic forms over F which become isotropic over K but no proper subforms of them will become isotropic over K . Knowing the K -minimal forms amounts to knowing which anisotropic forms over F become isotropic over K . This problem is related to the excellence problem and to some extent also to the problem of determining $W(K/F)$ as will become clear from our investigations. Thus, it is not surprising that the machinery we develop also leads to some extensions and generalizations of some of the results in the papers mentioned above, most notably [ELW 3].

It is well-known that if π is an anisotropic Pfister form over F then $W(F(\pi)/F) = \pi WF$ (see, e.g., [S, Ch.4, Theorem 5.4(iv)]). Thus, it seems natural to ask whether $W(K/F)$ is generated by $\{\rho_i\}$ where $K = F(\{\rho_i\})$ is a Pfister extension, i.e.,

$$W(F(\{\rho_i\})/F) = \sum_i \rho_i WF . \quad (1)$$

In [ELW 1], F was called *n-amenable* if equation (1) holds for any finite set $\{\rho_i\}$ of Pfister forms of fold $\leq n$, and F was said to be *amenable* if F is *n-amenable* for every $n \geq 1$.

In the context of our paper, it proves useful to extend this definition. We call F *co- n -amenable* if equation (1) holds for any finite set $\{\rho_i\}$ of Pfister forms of fold $\geq n$. Thus, F being amenable is the same as F being co-1-amenable. In analogy to the definition of amenability, we say that F is *n -Pfister excellent* (resp. *co- n -Pfister excellent*) if for every finite set $\{\rho_i\}$ of Pfister forms over F of fold $\leq n$ (resp. $\geq n$), the Pfister extension $F(\{\rho_i\})/F$ is excellent. We call F *Pfister excellent* if F is co-1-Pfister excellent (or, equivalently, if F is n -Pfister excellent for every n).

Recall that the Hasse number $\tilde{u}(F)$ of F is defined to be the supremum of the dimensions of all anisotropic forms over F which are indefinite with respect to each ordering on F (if there are any). If F is not formally real (i.e. there are no orderings on F), then $\tilde{u}(F)$ coincides with the u -invariant of F , where $u(F)$ (in the generalized sense) is defined to be the supremum of the dimensions of all anisotropic torsion forms over F . Let us now state a summary of known “positive” results. In parts (i)–(iv), ρ , σ , and π will denote anisotropic Pfister forms over F with $\dim \rho = 2$ and $\dim \sigma = 4$.

- (i) $W(F(\pi)/F) = \pi WF$ (see above);
- (ii) $F(\rho)/F$ and $F(\sigma)/F$ are excellent (see, e.g., [L1, Ch.VII, Lemma 3.1] or [S, Ch.2, Lemma 5.1] for $F(\rho)$, and [ELW 1, Appendix II by Arason] for $F(\sigma)$);
- (iii) If σ is universal then $F(\rho, \sigma)/F$ is excellent (see [ELW 1, Corollary 2.5]);
- (iv) $W(F(\rho, \pi)/F) = \rho WF + \pi WF$, $W(F(\sigma, \pi)/F) = \sigma WF + \pi WF$, and if σ is universal then $W(F(\rho, \sigma, \pi)/F) = \rho WF + \sigma WF + \pi WF$. (cf. [ELW 1, Corollary 2.12 and p. 466]);
- (v) If $\tilde{u}(F) \leq 2^n$ ($n = 1, 2$) then F is co- n -amenable and co- n -Pfister excellent (so F is amenable and Pfister excellent if $\tilde{u}(F) \leq 2$). Furthermore, if $\tilde{u}(F) = 4$ and F is 1-amenable then F is amenable and Pfister excellent (cf. [ELW 3, Main Theorem, Theorem 6.5]).

On the negative side, if ρ_1, ρ_2, ρ_3 are 1-fold Pfister forms then generally $F(\rho_1, \rho_2)/F$ is not excellent and $W(F(\rho_1, \rho_2, \rho_3)/F) \neq \sum \rho_i WF$ (in fact, all biquadratic extensions over F are excellent iff all triquadratic extensions yield the “expected” Witt kernel, but these equivalent conditions generally do not hold, cf. [ELTW]). Thus, fields are neither 1-amenable nor 1-Pfister excellent in general. Izhboldin [I] showed recently, that if π is an anisotropic n -fold Pfister form, $n \geq 3$, then there always exists a field extension E/F such that $E(\pi)/E$ is not excellent. In particular, function fields of n -fold Pfister forms, $n \geq 3$, are generally not excellent.

We will extend the above list of positive results. Recall that F is called linked if the quaternion algebras form a subgroup in the Brauer group of F . It is well-known that if F is linked then $\tilde{u}(F) \in \{0, 1, 2, 4, 8\}$, and all these values do occur (cf. [E]). Also, F is called n -linked if any two n -fold Pfister forms over F have a common $(n-1)$ -fold subform. F is linked iff F is 2-linked.

- (vi) Let $n \geq 1$ and let $\tilde{u}(F) \leq 2^n$. Then F is co- n -amenable and co- $(n+1)$ -Pfister excellent. If F is $(n-1)$ -amenable then F is amenable. If F is n -Pfister excellent then F is Pfister excellent.
- (vii) If $\tilde{u}(F) \leq 6$ or if F is linked, then F is co-3-Pfister excellent.
- (viii) Let $n \geq 2$ and let F be an n -linked field with $I_t^{n+1} F = 0$. Suppose that $I^n F = J_n F$. Then F is co- n -amenable.

These results, especially those in (vi) and (viii), are of interest mainly in the case of

formally real fields because only then do there exist anisotropic Pfister forms of dimension $> 2^n$ which can be taken to construct Pfister extensions. If one restricts oneself to function fields of n -fold Pfister forms in the above statements about amenability, then these results become a special case of the Linked Group Theorem which we state in the next section.

This paper is structured as follows. In the next section, we will recall several important results which we will need in our proofs. We will also say a little about (n -)linked fields and about fields with finite Hasse number. Some familiarity with the basic facts about SAP fields is assumed on the reader's part throughout this paper. The relevant facts can be found in [L2], [KS], and [ELP]. Many of the proofs consist of rather intricate arguments using signatures of quadratic forms, and very often we will use the fact that certain sets of orderings we define are clopen sets and that F is SAP (which is always guaranteed if F is n -linked or the Hasse number is finite). We will not always prove these facts explicitly since they can be readily checked by the reader. In § 3, we provide two main lemmas which are somewhat technical in nature but designed in a way so that they can be readily used in the proofs of the results mentioned above and in our investigations of K -minimal forms. We will prove above results in § 4. In § 5, we provide bounds for K -minimal forms where K/F is a Pfister extension over a field with finite Hasse number. For some of the Pfister extensions K/F we will be dealing with, we will give a characterization of the K -minimal forms in § 6. To round things off, we explicitly compute the K -minimal forms for certain Pfister extensions K/F in § 7 to demonstrate the power of our approach. In one example, the field F will be a formally real global field, and in the other example F will be $\mathbf{R}(t)$, the rational function field in one variable over the real numbers.

2 Preliminaries

By $I^n F$ we denote the n -th power of the fundamental ideal IF of even-dimensional forms in WF . $I^n F$ is additively generated by the n -fold Pfister forms $\langle\langle a_1, \dots, a_n \rangle\rangle = \langle 1, a_1 \rangle \otimes \dots \otimes \langle 1, a_n \rangle$. The set of forms over F isometric (resp. similar) to n -fold Pfister forms will be denoted by $P_n F$ (resp. $GP_n F$). PF denotes the set of all Pfister forms over F (GPF is defined accordingly). $W_t F$ denotes the torsion part of the Witt ring WF , and we put $I_t^n F = I^n F \cap W_t F$. $J_n F$ denotes the ideal of forms of degree $\geq n$ in WF as defined by Knebusch in [K1]. It is conjectured that $I^n F = J_n F$ for all n and all fields F , and it is known to be true in general when $n \leq 4$ (and also for $n = 5$ due to some unpublished results by Rost, see [Ka, Théorème 2.8] and the remarks and references there; recently, Voevodsky announced certain results which would imply that this is true for all n). A generic splitting tower of a form φ over F is, in our context, a tower of fields $F = F_0 \subset F_1 \subset \dots \subset F_h$ with the F_i defined as follows. Let $\varphi_0 \simeq \varphi_{\text{an}}$, and for $i \geq 1$ let $F_i = F_{i-1}(\varphi_{i-1})$ and $\varphi_i \simeq ((\varphi_{i-1})_{F_i})_{\text{an}}$. $h \geq 0$ is the smallest integer for which $\dim \varphi_h \leq 1$ and F_h is called the generic splitting field of φ (cf. [K1] for this construction and other facts about generic splitting towers).

Let $\Phi \subset P_n F$ be a set of n -fold Pfister forms over F . Following the terminology in [ELW3], we call Φ a linked group of n -fold Pfister forms if $\overline{\Phi} = \{\pi \pmod{I^{n+1}F} \mid \pi \in \Phi\}$ is a subgroup of $I^n F / I^{n+1} F$. In other words, Φ is a linked group of n -fold Pfister forms if to any $\pi_1, \pi_2 \in \Phi$ there exists $\pi_3 \in \Phi$ such that $\pi_1 - \pi_2 \equiv \pi_3 \pmod{I^{n+1}F}$. Furthermore, we say that $\Phi \subset P_n F$ generates a linked group of Pfister forms if there exists a linked group $\Psi \subset P_n F$ such that $\overline{\Phi}$ generates $\overline{\Psi}$. Recall that two n -fold Pfister forms π_1, π_2 are called linked if there exist an $(n-1)$ -fold Pfister form ρ and 1-fold Pfister forms τ_1, τ_2 such that $\pi_i \simeq \rho \otimes \tau_i$, $i = 1, 2$. In a linked group of n -fold Pfister forms any two Pfister forms

are linked. The field F is called n -linked if $P_n F$ is a linked group of Pfister forms. If $n \geq 2$ then it is easy to see that F being n -linked implies that F is m -linked for all $m \geq n$. If $n = 2$ then F is simply called linked.

The next few results show how certain properties of F imply that F is n -linked. Here and throughout the remainder of the paper, we denote the space of orderings on F by X_F , and the signature of a quadratic form φ with respect to $P \in X_F$ will be denoted by $\text{sgn}_P \varphi$. The next lemma is rather obvious and we omit its proof.

Lemma 2.1 *Let φ be a form over F . Then*

$$\dim \varphi_{\text{an}} \leq \sup\{\tilde{u}(F), |\text{sgn}_P \varphi|; P \in X_F\}.$$

Lemma 2.2 *F is n -linked if one of the following properties holds:*

- (i) $\tilde{u}(F) < 2^n + 2^{n-1}$.
- (ii) F is SAP and $I_t^n F = 0$.

Proof. (i) Let $\pi_1, \pi_2 \in P_n F$. If F is formally real then, for all $P \in X_F$, we have $\text{sgn}_P \pi_i \in \{0, 2^n\}$ and thus $|\text{sgn}_P(\pi_1 \perp -\pi_2)| \in \{0, 2^n\}$. In any case, the previous lemma implies that $\dim(\pi_1 \perp -\pi_2)_{\text{an}} < 2^n + 2^{n-1}$. Hence, by [EL 1, Theorem 4.5], $\dim(\pi_1 \perp -\pi_2)_{\text{an}} = 0$ or 2^n and π_1 and π_2 are linked.

(ii) If F is not formally real then there are no anisotropic n -fold Pfister forms and there is nothing to show. So assume that F is formally real and that $\pi_1, \pi_2 \in P_n F$. As F is SAP and as $|\text{sgn}_P(\pi_1 \perp -\pi_2)| \in \{0, 2^n\}$ there exist $\pi \in P_n F$ and $x \in \dot{F}$ such that

$$\text{sgn}_P \pi = |\text{sgn}_P(\pi_1 \perp -\pi_2)| \text{ for all } P \in X_F$$

and

$$\begin{aligned} x <_P 0 & \text{ for all } P \in X_F \text{ with } \text{sgn}_P(\pi_1 \perp -\pi_2) = -2^n \\ x >_P 0 & \text{ otherwise.} \end{aligned}$$

Thus, $\text{sgn}_P(\pi_1 \perp -\pi_2) = \text{sgn}_P x\pi$ for all $P \in X_F$ and therefore

$$\pi_1 \perp -\pi_2 \perp -x\pi \in I_t^n F = 0.$$

Hence, $\pi_1 \perp -\pi_2 = x\pi$ in WF which shows that π_1 and π_2 are linked. \square

Example 2.3 Generally, it is not true that F being SAP and $I_t^n F = 0$ implies that F is $(n-1)$ -linked. For let F be a field with $\tilde{u}(F) = 6$ (such fields exist by Merkurjev's results on the u -invariant of fields [M]). Clearly, F is SAP as $\tilde{u}(F)$ is finite, and $I_t^3 F = 0$ by the Arason-Pfister Hauptsatz (see below). However, F is not (2-)linked as the Hasse number of a linked field is never 6 (see the remarks in the introduction). As another example, consider the Laurent series field in one variable over the rational numbers, $F = \mathbf{Q}((t))$. Now $u(\mathbf{Q}) = 4$, and thus, by Springer's theorem (cf. [S, Ch. 6, Corollary 2.6]), $u(F) = 8$. Hence, $I_t^4 F = 0$. On the other hand, F is SAP (cf. [L 2, p. 124]). However, F is not 3-linked because, for example, $\langle\langle 1, 1, 1 \rangle\rangle$ and $\langle\langle 1, -3, t \rangle\rangle$ are not linked as one can easily check. It will turn out that F is co-4-amenable (see Example 4.3).

For reference reasons, let us now state some well-known and important results which we will use frequently.

Proposition 2.4 (i) (Cassels-Pfister subform theorem, cf. [S, Ch.4, Theorem 5.4].) *Let φ be an anisotropic form over F and ψ be another form over F . If $\varphi_{F(\psi)}$ is hyperbolic then $a\psi \subset \varphi$ for any $a \in D(\varphi)D(\psi)$. In particular, $\dim \varphi \geq \dim \psi$.*

(ii) (Arason-Pfister Hauptsatz (APH), cf. [S, Ch.4, Theorem 5.6].) *Let φ be an anisotropic form in $I^n F$. Then $\dim \varphi \geq 2^n$. Furthermore, if $\dim \varphi = 2^n$ then φ is similar to an n -fold Pfister form.*

(iii) (“Anisotropy Criterion”, [H4, Theorem 1].) *Let φ and ψ be forms over F and let φ be anisotropic. If $\dim \varphi \leq 2^n < \dim \psi$ for some n then $\varphi_{F(\psi)}$ is anisotropic.*

(iv) (Extension of orderings, cf. [ELW 2, Theorem 3.5].) *Let F be formally real and let φ be a form over F . Then $P \in X_F$ extends to an ordering on $F(\varphi)$ if and only if φ is indefinite with respect to P , i.e., $|\operatorname{sgn}_P \varphi| < \dim \varphi$. In particular, if K is a generic splitting field as defined above then $P \in X_F$ extends to an ordering on K if and only if $|\operatorname{sgn}_P \varphi| \leq 1$.*

Here, $a\psi \subset \varphi$ means that $a\psi$ is a subform of φ , i.e., there exists $\tau \in WF$ such that $\varphi \simeq a\psi \perp \tau$. We will use this notation frequently. Also, $D(\varphi) = \{x \in \dot{F} \mid \langle x \rangle \subset \varphi\}$ denotes the set of non-zero elements in F which are represented by φ .

The last result we shall mention will play a most crucial role in our investigations. It is the Linked Group Theorem (we will refer to it as LGT) and was proved in [ELW 3, Theorem 2.1].

Theorem 2.5 (Linked Group Theorem.) *Let $\{\rho_i\} \subset P_n F$ generate a linked group Φ of Pfister forms over F . Let E/F be a field extension and let $L = E(\{\rho_i\})$. Then*

$$\Phi \subseteq \sum_i \rho_i W E \subseteq W(L/E).$$

Moreover, if $\psi \in P_n E$ then $\psi \in W(L/E)$ if and only if there exists $\rho \in \Phi$ such that $\psi \simeq \rho_E$.

Our version here is a touch more general than the original one in which only the case $E = F$ was considered. However, only very minor modifications of the original proof are needed to yield the present form of the theorem.

3 Two lemmas

Lemma 3.1 *Let F be formally real and n -linked. Let $Y \subset X_F$ be clopen and let E/F be a field extension such that each $P \in Y$ extends to E . Let $\varphi \in WF$ such that $\varphi_E \in P_n E$. Let $\{\pi_\alpha\} \subset PF$ and suppose that $\varphi_{E(\{\pi_\alpha\})} = 0$. Then there exists $\pi \in P_n F \cap \sum (\pi_\alpha WF \cap GP_n F)$ such that $\operatorname{sgn}_P \varphi = \operatorname{sgn}_P \pi$ for all $P \in Y$.*

Proof. If φ_E is hyperbolic then $\operatorname{sgn}_P \varphi = 0$ for all $P \in Y$ and we can choose $\pi \in P_n F$ to be hyperbolic. So suppose φ_E is anisotropic. Now $\dim \varphi = 2^n$ and $\varphi_{E(\{\pi_\alpha\})} = 0$. Thus, we may assume that $\{\pi_\alpha\}$ is finite and all π_α are of fold $\leq n$. So let $\{\pi_\alpha\} = \{\rho_1, \dots, \rho_r, \sigma_1, \dots, \sigma_s\}$ with $r + s > 0$, $\rho_i \in PF$ of fold $\leq n - 1$, and $\sigma_j \in P_n F$. We define a tower of fields $E = L_0 \subseteq L_1 \subseteq \dots \subseteq L_r = L \subseteq M$ by $L_i = L_{i-1}(\rho_i)$ for $1 \leq i \leq r$ and $M = L(\sigma_1, \dots, \sigma_s)$.

Now $\varphi_L \in W(M/L) \cap P_n L$. Since F is n -linked, LGT implies that there exists $\sigma \in P_n F \cap \sum_{i=1}^s \sigma_i WF$ with $\varphi_L \simeq \sigma_L$. It is clear by our assumptions that $\operatorname{sgn}_P \varphi, \operatorname{sgn}_P \sigma \in \{0, 2^n\}$ for all $P \in Y$. Let $\psi \simeq \varphi \perp -\sigma$ over F . Then $|\operatorname{sgn}_P \psi| \in \{0, 2^n\}$ for all $P \in Y$ as well. Since F is SAP (recall that F is n -linked!), we can choose $x \in \dot{F}$ such that

$\text{sgn}_P x\psi \in \{0, 2^n\}$ for all $P \in Y$. Note also that $\psi_E \in W(L/E)$. Let now $Z = \{P \in Y \mid \text{sgn}_P x\psi = 2^n\}$, $Y_0 = \emptyset$, and for $1 \leq i \leq r$ let

$$Y_i = \{P \in Z \setminus (Y_0 \cup \dots \cup Y_{i-1}) \mid \rho_i \text{ is definite at } P\}.$$

Then $P \in Z$ extends to L_{i-1} but not to L_i iff $P \in Y_i$. Since $\psi_L = 0$, no $P \in Z$ can extend to L . Hence, Z is the disjoint union of the Y_i 's. Let now $t_i \in \bar{F}$ and $\tau_i \simeq \langle\langle 1, \dots, 1, t_i \rangle\rangle \in PF$ be such that $\tau_i \otimes \rho_i \in P_n F$, τ_i is definite for all $P \in Y_i$ and torsion for all $P \in X_F \setminus Y_i$. Such t_i 's exist as F is SAP and the Y_i are clopen sets. One readily checks that $\text{sgn}_P \tau_i \otimes \rho_i = 2^n$ for all $P \in Y_i$ and 0 for all $P \in X_F \setminus Y_i$. Hence,

$$\text{sgn}_P \sum_{i=1}^r \tau_i \otimes \rho_i = \begin{cases} 2^n & \text{for } P \in Z \\ 0 & \text{for } P \in X_F \setminus Z \end{cases}$$

and thus $\text{sgn}_P \sum_{i=1}^r \tau_i \otimes \rho_i = \text{sgn}_P x\psi$ for all $P \in Y$. Hence,

$$\text{sgn}_P \varphi = \text{sgn}_P (\sigma + \sum_{i=1}^r x\tau_i \otimes \rho_i) \text{ for all } P \in Y.$$

Since F is n -linked, there exists $\pi \in P_n F$ with $\sigma + \sum_{i=1}^r x\tau_i \otimes \rho_i \equiv \pi \pmod{I^{n+1}F}$. In particular, this yields $\text{sgn}_P \varphi \equiv \text{sgn}_P \pi \pmod{2^{n+1}}$ for all $P \in Y$. In fact, $\text{sgn}_P \varphi = \text{sgn}_P \pi$ for all $P \in Y$ because $\text{sgn}_P \varphi, \text{sgn}_P \pi \in \{0, 2^n\}$. It is now easy to conclude, using that F is n -linked, that $\pi \in \sigma WF + \sum (\tau_i \otimes \rho_i) WF$. Moreover, n -linkage readily implies that

$$\pi = \sum_{i=1}^r x_i \tau_i \otimes \rho_i + \sum_{j=1}^s y_j \sigma_j \in \sum (\pi_\alpha WF \cap GP_n F)$$

for suitable $x_i, y_j \in \bar{F}$. \square

Lemma 3.2 *Let F be formally real with $\tilde{u}(F) < 2^n$ for some $n \geq 1$. Let $\varphi \in WF$ be anisotropic with $\max\{\tilde{u}(F), 2^{n-1}\} < \dim \varphi \leq 2^n$. Let $\{\pi_\alpha\} \subset PF$ be a set of Pfister forms of fold ≥ 1 and $K = F(\{\pi_\alpha\})$. Suppose that φ_K is isotropic. Then there exists an n -fold Pfister form $\pi \in \sum (\pi_\alpha WF \cap GP_n F)$ such that φ contains a Pfister neighbor of π . In particular, there exists $\psi \in WF$ with $\dim \psi < \dim \varphi$ such that $\varphi_K = \psi_K$ in WK .*

Proof. Since $\tilde{u}(F) < 2^n$ we have that F is n -linked and SAP. Now let $0 \leq m < 2^{n-1}$ such that $\dim \varphi = 2^n - m$. Since F is SAP, we may assume that after scaling we have $\text{sgn}_P \varphi \geq 0$ for all $P \in X_F$. Let

$$\begin{aligned} Y_0 &= \{P \in X_F \mid \text{sgn}_P \varphi = \dim \varphi, \text{ i.e., } \varphi \text{ is definite at } P\} \\ Y_1 &= \{P \in X_F \mid 0 \leq \text{sgn}_P \varphi \leq m\} \end{aligned}$$

As F is SAP, there exists $\varphi' \in WF$, $\dim \varphi' = m$, such that $\text{sgn}_P \varphi' = m$ (resp. $\text{sgn}_P \varphi' = -\text{sgn}_P \varphi$) for all $P \in Y_0$ (resp. $P \in Y_1$). Let $\gamma \simeq \varphi \perp \varphi'$. Then $\dim \gamma = 2^n$ and we have

$$\text{sgn}_P \gamma = \begin{cases} 2^n & \text{for all } P \in Y_0 \\ 0 & \text{for all } P \in Y_1 \end{cases}$$

Again by the SAP property, there exists $\tau \in P_n F$ such that $\text{sgn}_P \tau = \text{sgn}_P \gamma$ for all $P \in Y_0 \cup Y_1 = Y$. Let E be the generic splitting field of $\tau \perp -\gamma$. Since $\tau \perp -\gamma$ has signature 0 at each $P \in Y_0 \cup Y_1 = Y$, it follows that each $P \in Y$ extends to E (see

Proposition 2.4(iv)), and we also have $\tau_E \simeq \gamma_E \in P_n E$. Let $L = E(\{\pi_\alpha\})$. Now φ_K and hence γ_K is isotropic. Thus, $\gamma_L \simeq \tau_L \in P_n L$ is hyperbolic as $K \subset L$. By Lemma 3.1, there exists $\pi \in \sum(\pi_\alpha W F \cap G P_n F)$ such that $\text{sgn}_P \gamma = \text{sgn}_P \pi$ for all $P \in Y$. Hence, using that $m < \text{sgn}_P \varphi < 2^n - m$ for all $P \in X_F \setminus Y$ and that $\text{sgn}_P \pi \in \{0, 2^n\}$ for all P , we get

$$|\text{sgn}_P(\varphi \perp -\pi)| \leq \begin{cases} m & \text{for all } P \in Y \\ 2^n - m - 1 & \text{otherwise.} \end{cases}$$

Now $m < 2^n - m = \dim \varphi$ and $\tilde{u}(F) < \dim \varphi$ and we get by Lemma 2.1

$$\dim(\varphi \perp -\pi)_{\text{an}} \leq \max\{2^n - m - 1, \tilde{u}(F)\} < \dim \varphi .$$

Hence, $i_W(\varphi \perp -\pi) > \frac{1}{2} \dim \pi$ which implies that φ contains a subform of π of dimension $\geq 2^{n-1} + 1$, i.e., φ contains a Pfister neighbor of π .

The remaining statement of the lemma follows readily by putting $\psi \simeq (\varphi \perp -\pi)_{\text{an}}$ because $\pi_K = 0$. \square

4 Amenability and excellence results

Theorem 4.1 *Let $n \geq 2$. Let F be an n -linked field with $I_i^{n+1} F = 0$. Suppose that $I^n F = J_n F$. Then F is co- n -amenable.*

Proof. Let $\Pi \subset \bigcup_{r \geq n} P_r F$, i.e., Π is a set of Pfister forms of fold $\geq n$ over F . Let $\mathfrak{S} = \sum_{\pi \in \Pi} \pi W F$. For $m \geq n$ let $\Pi_m = \{\pi \mid \pi \in \bigcup_{n \leq r \leq m} P_r F \cap \Pi\}$, i.e. Π_m consists of all Pfister forms in Π of fold $\leq m$. Let $\mathfrak{S}_m = \sum_{\pi \in \Pi_m} \pi W F$. We obviously have $\mathfrak{S} = \bigcup_{m \geq n} \mathfrak{S}_m$. Let $K = F(\Pi)$. We have to show that $W(K/F) = \mathfrak{S}$.

Clearly, $\mathfrak{S} \subset W(K/F)$. So let $\psi \in W(K/F)$ be anisotropic. Since K is obtained by taking the function field of forms of dimension $\geq 2^n$ and since ψ becomes hyperbolic over K , it follows from [K1, Proposition 6.11] that the degree of ψ has to be $\geq n$, i.e. $\psi \in J_n F$. By assumption, we then have that $\psi \in I^n F$. By [E, Lemma 4.4], there are forms $\alpha_i \in P_i F$ and $z_i \in F$, $n \leq i \leq r$ for some $r \geq n$, such that

$$\psi = \sum_{i=n}^r z_i \alpha_i \quad \text{in } W F.$$

It follows immediately from [ELW1, Lemma 1.3] that $\alpha_i \in W(K/F)$ for $n \leq i \leq r$. However, as $\dim \alpha_i = 2^i$ we must already have that α_i becomes hyperbolic over $F(\Pi_i)$. Now for $i = n$ we know that Π_n generates a linked group of Pfister forms because F is n -linked. It readily follows from LGT that $\alpha_n \in \mathfrak{S}_n$.

Now if all $\alpha_i = 0$ in $W F$ for $i > n$ then we are done. So suppose $i > n$ and $\alpha_i \neq 0$. Then $\alpha_i \in P_i F$ is anisotropic, and because $I_i^i F = 0$ by assumption we necessarily have that F is formally real. Also, F is i -linked because it is n -linked. Thus, we can apply Lemma 3.1 to conclude that there exists a form $\beta \in \mathfrak{S}_i \cap P_i F$ such that $\text{sgn}_P \alpha_i = \text{sgn}_P \beta$ for all $P \in X_F$. Now the form $\alpha_i \perp -\beta$ is in $I^i F$ because $\alpha_i, \beta \in I^i F$, and it has signature 0 at each ordering of F . Hence, $\alpha_i \perp -\beta \in I_i^i F = 0$. It follows that $\alpha_i = \beta$ in $W F$ and thus $\alpha_i \in \mathfrak{S}_i$. Therefore,

$$\psi = \sum_{i=n}^r z_i \alpha_i \in \sum_{i=n}^r \mathfrak{S}_i \subset \mathfrak{S} .$$

\square

Corollary 4.2 *Let $n \geq 2$ and suppose that $I^n F = J_n F$. Then F is co- n -amenable if F satisfies one of the following conditions:*

- (i) F is $(n - 1)$ -linked.
- (ii) $\tilde{u}(F) < 2^n + 2^{n-1}$.
- (iii) F is SAP and $I_t^n F = 0$.
- (iv) F is SAP and $u(F) < 2^n$.
- (v) F is n -linked and $u(F) < 2^{n+1}$.

In particular, if F satisfies any of these conditions and if $n \leq 4$ then F is co- n -amenable.

Proof. In order to apply the previous theorem we have to verify that F is n -linked and $I_t^{n+1} F = 0$. In (i), this follows from [EL 2, Lemma 2.3, Cor. 2.8]. It is clear that in the remaining cases $I_t^{n+1} F = 0$. F is n -linked in (ii) and (iii) by Lemma 2.2. One readily sees that (iv) implies (iii). F is n -linked in (v) by assumption. Finally, if $n \leq 4$ then, as mentioned in the introduction, one has $I^n F = J_n F$ which then implies the last statement. \square

It is well-known that if $\tilde{u}(F) \leq 2^n$ then $I^n F = J_n F$ (see, e.g., [AK, Lemma 2], or [F, Lemma 4.8]). The above then implies that if $\tilde{u}(F) \leq 2^n$ then F is co- n -amenable. We do not state this here as another corollary because below we will prove stronger results under the assumption that $\tilde{u}(F) \leq 2^n$ (Theorem 4.4 and corollary).

Example 4.3 Let $F = \mathbf{Q}(\langle t \rangle)$ be the Laurent series field in one variable over the rational numbers. F is easily seen to be 4-linked, the only anisotropic 4-fold Pfister forms being $\langle\langle 1, 1, 1, 1 \rangle\rangle$ and $\langle\langle 1, 1, 1, \pm t \rangle\rangle$. Also, by Springer's theorem, $u(F) = 2u(\mathbf{Q}) = 8$. By Corollary 4.2, F is co-4-amenable. Note that $\tilde{u}(F) = \infty$ as for example the form $\langle 1, -2 \rangle \perp (n \times \langle t \rangle)$ is t.i. and anisotropic for each n .

Before we state our next main result, we will fix some notation. Let $\{\sigma_i\} \subset PF$ be a set of Pfister forms of fold $\leq n - 1$, let $\{\rho_j\} \subset PF$ be a set of Pfister forms of fold n , and let $\{\pi_k\} \subset PF$ be a set of Pfister forms of fold $\geq n + 1$. Let

$$\begin{aligned} K_1 &= F(\{\sigma_i\}) \\ K_2 &= F(\{\sigma_i\}, \{\rho_j\}) \\ K_3 &= F(\{\sigma_i\}, \{\rho_j\}, \{\pi_k\}) \end{aligned}$$

and let

$$\begin{aligned} \mathfrak{S}_1 &= \sum_i \sigma_i WF \\ \mathfrak{S}_2 &= \sum_i \sigma_i WF + \sum_j \rho_j WF \\ \mathfrak{S}_3 &= \sum_i \sigma_i WF + \sum_j \rho_j WF + \sum_k \pi_k WF . \end{aligned}$$

Clearly, $K_1 \subset K_2 \subset K_3$, $\mathfrak{S}_1 \subset \mathfrak{S}_2 \subset \mathfrak{S}_3$, and $\mathfrak{S}_m \subset W(K_m/F)$ for $1 \leq m \leq 3$.

Theorem 4.4 *Suppose $\tilde{u}(F) \leq 2^n$.*

- (i) *If K_2/F is excellent then K_3/F is excellent.*
- (ii) *If $W(K_1/F) = \mathfrak{S}_1$ then $W(K_3/F) = \mathfrak{S}_3$.*

Proof. (i) We have to show that if $\varphi \in WF$ is anisotropic and φ_{K_3} is isotropic then $(\varphi_{K_3})_{\text{an}}$ is defined over F .

If $\dim \varphi \leq 2^n$, then by the Anisotropy Criterion φ must become isotropic already over K_2 . As K_2/F is excellent there exists an anisotropic $\mu \in WF$ with $\mu_{K_2} \simeq (\varphi_{K_2})_{\text{an}}$. It

follows again from the Anisotropy Criterion that μ_{K_2} stays anisotropic over K_3 . Therefore, $\mu_{K_3} \simeq (\varphi_{K_3})_{\text{an}}$ and we are done.

Now if $\dim \varphi > 2^n$ then by Lemma 3.2 there exists $\psi \in WF$ with $\dim \psi < \dim \varphi$ such that $\varphi_{K_3} = \psi_{K_3}$ in WK_3 . The assertion now follows easily by using induction on $\dim \varphi$.

(ii) We want to show that $W(K_3/F) = \mathfrak{S}_3$. So let $\varphi \in W(K_3/F)$ be anisotropic. If $\dim \varphi < 2^n$ we already must have $\varphi \in W(K_1/F) = \mathfrak{S}_1 \subset \mathfrak{S}_3$ because anisotropic forms of dimension $< 2^n$ cannot become hyperbolic over the function field of forms of dimension $\geq 2^n$ by the Cassels-Pfister subform theorem. So we are done in this case.

So suppose $\dim \varphi = 2^n$. If $\dim(\varphi_{K_1})_{\text{an}} < 2^n$ we already have $\varphi_{K_1} = 0$ by the same argument as above. Again, $\varphi \in W(K_1/F) = \mathfrak{S}_1 \subset \mathfrak{S}_3$. So suppose $\dim(\varphi_{K_1})_{\text{an}} = \dim \varphi = 2^n$. As K_3/K_1 is obtained by taking function fields of forms of dimension $\geq 2^n$, we must have $\deg \varphi_{K_1} \geq n$ because $\varphi_{K_3} = 0$ (see [K1, Proposition 6.11]). But $\dim \varphi_{K_1} = 2^n$ and thus $\varphi_{K_1} \in GP_n K_1$. After possibly scaling, we may assume that φ represents 1 over F and thus over K_1 . Hence, $\varphi_{K_1} \in P_n K_1$. Now φ_{K_3} is hyperbolic. It is immediate that therefore φ_{K_2} is already hyperbolic as K_3 is a function field extension of forms of dimension $\geq 2^{n+1}$ over K_2 . Hence,

$$\varphi_{K_1} \in W(K_2/K_1) \cap P_n K_1 .$$

Now $K_2 = K_1(\{\rho_j\})$, and as F is n -linked because $\tilde{u}(F) \leq 2^n$, we have that $\{\rho_j\}$ generates a linked group on n -fold Pfister forms over F . Thus, by LGT, there exists $\pi \in P_n F \cap (\sum \rho_j WF)$ such that $\pi_{K_1} \simeq \varphi_{K_1}$. In particular,

$$\varphi \perp -\pi \in W(K_1/F) = \mathfrak{S}_1 .$$

Thus, $\varphi \in \mathfrak{S}_1 + \sum \rho_j WF \subset \mathfrak{S}_3$.

Now if $\dim \varphi > 2^n$ and φ_{K_3} is hyperbolic then, by Lemma 3.2, there exists $\pi \in \mathfrak{S}_3 \cap GPF$ such that for $\psi \simeq (\varphi \perp -\pi)_{\text{an}}$ we have $\dim \psi < \dim \varphi$. As $\pi \in \mathfrak{S}_3 \subset W(K_3/F)$ we have $\psi_{K_3} = \varphi_{K_3} \perp -\pi_{K_3} = 0$ in WK_3 . Using induction on dimension we conclude that $\psi \in \mathfrak{S}_3$ and thus $\varphi \in \mathfrak{S}_3$ because $\varphi = \psi \perp \pi$ in WF . \square

Corollary 4.5 *Let $\tilde{u}(F) \leq 2^n$. Then the following holds.*

- (i) *F is co- $(n+1)$ -Pfister excellent. If F is n -Pfister excellent then F is Pfister excellent.*
- (ii) *F is co- n -amenable. If F is $(n-1)$ -amenable then F is amenable.*

We can improve on the above results if we assume that the Hasse number is small enough. These improvements are achieved by invoking well-understood properties of forms of small dimension. In particular, we have the following theorem.

Theorem 4.6 (i) *If $\tilde{u}(F) \leq 2$ then F is Pfister excellent.*

- (ii) *If $\tilde{u}(F) \leq 4$ then F is co-2-Pfister excellent.*
- (iii) *If $\tilde{u}(F) \leq 4$ and F is 1-amenable then F is Pfister excellent.*
- (iv) *If $\tilde{u}(F) \leq 6$ then F is co-3-Pfister excellent.*
- (v) *If F is linked then F is co-3-Pfister excellent.*

Our proof of parts (i)–(iii) will be shorter than the one given in [ELW 3] although some of our arguments will be in a similar spirit. To prove parts (iv) and (v), we will need the following result.

Proposition 4.7 *Let $\{\rho_i\} \subset P_3F$ generate a linked group Φ of 3-fold Pfister forms over F and let $K = F(\{\rho_i\})$. Let $\varphi \in WF$ be anisotropic and suppose that one of the following conditions is satisfied.*

- (i) $\dim \varphi \leq 6$;
- (ii) $\dim \varphi = 7$, or $\dim \varphi = 8$ and $\varphi \in I^2F$, and in addition $c(\varphi) \in BrF$ can be represented by a single quaternion algebra, i.e., $c(\varphi) = (x, y)_F$ in BrF for some $x, y \in \dot{F}$;
- (iii) F is linked and $\dim \varphi \leq 8$.

Then φ_K is isotropic if and only if there exists $\pi \in \Phi$ such that φ contains a Pfister neighbor of π .

Proof. It clearly suffices to show the “only if”-part. By the Anisotropy Criterion we must have $\dim \varphi \geq 5$ for φ_K to be isotropic. If $\dim \varphi \leq 6$ the above statement was essentially shown in [H 3, Corollary 4.2] in the case of a function field of a single 3-fold Pfister form. The above more general statement can be shown in a similar way after invoking LGT and, at a certain point, the fact that anisotropic 6-dimensional forms in I^2 stay anisotropic over function fields of forms of dimension ≥ 7 (see, e.g., [L 3, Theorem 4.3]). We leave it to the reader to fill in the details.

(ii) has been shown in [H 5, Theorem 4.5, Corollary 4.6].

Finally suppose we are in the situation of (iii) and let us assume that φ_K is isotropic. If $\dim \varphi \leq 6$ we are done by part (i). So suppose $\dim \varphi = 7$ or 8. As F is linked there exist $x, y \in \dot{F}$ such that $c(\varphi) = (x, y)_F \in BrF$. If $\dim \varphi = 7$, or if $\dim \varphi = 8$ and $\varphi \in I^2F$ the result then follows from (ii). So let us assume that $\dim \varphi = 8$ and that $d = d_{\pm}\varphi \neq 1 \in \dot{F}/\dot{F}^2$. Let $E = F(\sqrt{d})$. Suppose that φ_E is isotropic. Then there exist $z \in \dot{F}$ and $\alpha \in WF$ such that $\varphi \simeq \alpha \perp z\langle 1, -d \rangle$. This yields $d_{\pm}\alpha = 1$, i.e., α is a 6-dimensional form in I^2F . But over a linked field 6-dimensional I^2 -forms are always isotropic, a contradiction to φ being anisotropic. Hence, φ_E is anisotropic.

Let $L = E(\langle\langle -x, -y \rangle\rangle)$. If $\langle\langle -x, -y \rangle\rangle_E$ is isotropic then L/E is purely transcendental and φ stays anisotropic over L . If $\langle\langle -x, -y \rangle\rangle_E$ is anisotropic then obviously $\langle\langle -x, -y \rangle\rangle_L = 0$ or, equivalently, $(x, y)_L = 1$ in BrL . Thus, $\varphi_L \in I^3L$, and as $\dim \varphi = 8$ we have $\varphi_L \in GP_3L$. Suppose φ_L is isotropic. Then in fact $\varphi_L = 0$, i.e., $\varphi_E \in W(L/E)$ and there exists a form $\tau \in WE$, $\dim \tau = 2$, such that $\varphi_E \simeq \tau \otimes \langle\langle -x, -y \rangle\rangle_E$. In particular, $\varphi_E \in GP_3E$ and thus $c(\varphi_E) = 1 = (x, y)_E$ in BrE , which in turn implies that $\langle\langle -x, -y \rangle\rangle_E = 0$, a contradiction. Hence, $\varphi_L \in GP_3L$ is anisotropic. We may assume that φ represents 1 over F and thus, $\varphi_L \in P_3L$. Now clearly φ_L becomes isotropic and hence hyperbolic over $L(\{\rho_i\})$. By LGT, there exists $\pi \in \Phi$ such that $\varphi_L \simeq \pi_L$. As F is linked we have $\tilde{u}(F) \leq 8$ (cf. [E, Theorem 4.7]). As F is SAP there exists $a \in \dot{F}$ such that $|\text{sgn}_P(\varphi \perp -a\pi)| \leq 8$ for all $P \in X_F$. Let $\psi \simeq (\varphi \perp -a\pi)_{\text{an}}$. We then have $\dim \psi \leq \max\{\tilde{u}(F), |\text{sgn}_P(\varphi \perp -a\pi)|; P \in X_F\} \leq 8$.

Suppose $\dim \psi = 8$. We have $d_{\pm}\psi = d_{\pm}(\varphi \perp -\pi) = d_{\pm}\varphi = d$ and $c(\psi) = c(\varphi \perp -\pi) = c(\varphi) = (x, y)_F$. By the same argument as before, we see that ψ_L is anisotropic. But

$$\psi_L = \varphi_L \perp -a\pi_L = \pi_L \perp -a\pi_L \quad \text{in } WL.$$

Comparing dimensions, we see that $\pi_L \perp -a\pi_L \in P_4L$ is isotropic and hence hyperbolic, which in turn implies that $\psi_L = 0$ in WL , a contradiction. Hence, $\dim \psi < 8$, i.e., $\dim \psi \leq 6$. Then we have $i_W(\varphi \perp -a\pi) \geq 5$ and there exists $\chi \in WF$ with $\dim \chi = 5$, $\chi \subset \varphi$ and $\chi \subset a\pi$, which concludes the proof. \square

An analogous result to that in (ii) also holds if $\varphi \in I^2F$, $\dim \varphi = 8$, and $c(\varphi) = (x, y)_F \otimes (u, v)_F$ in BrF , where $x, y, u, v \in \dot{F}$ (i.e., $c(\varphi)$ can be represented by a bi-quaternion algebra in BrF , or equivalently, $\text{ind } c(\varphi) \leq 4$). This follows easily from [Lag, Théorème 2(iv)] after a minor modification of the proof given there. If $\varphi \in I^2F$ and $\dim \varphi = 8$ then one easily shows that $\text{ind } c(\varphi) \leq 8$, and if it is 8 then φ is necessarily anisotropic and it follows from Merkurjev's index reduction theorem [M] that φ will stay anisotropic over function fields of 3-fold Pfister forms. If $\dim \varphi = 8$ and $d_{\pm}\varphi \neq 1$, then the above conclusion generally fails to be true as follows from Izhboldin's results in [I], see also [H 6].

Proof of Theorem 4.6. Let $n = 1$ in (i), $n = 2$ in (ii) and (iii), and $n = 3$ in (iv) and (v). We then have $\tilde{u}(F) \leq 2^n$. By the definition of (co- n) Pfister excellence and after invoking Theorem 4.4 it suffices to show that if $\{\rho_i\} \subset P_n F$ (resp. $\{\rho_i\} \subset P_1 F \cup P_2 F$ in part (iii)) and if $K = F(\{\rho_i\})$ then K/F is excellent. To show this, let $\varphi \in WF$ be anisotropic and suppose that φ_K is isotropic. If $\dim \varphi > \tilde{u}(F)$ then, by Lemma 3.2, there exists $\psi \in WF$ with $\dim \psi < \dim \varphi$ such that $\psi_K = \varphi_K$ in WK . Hence, it suffices to show that if $\varphi \in WF$ is anisotropic with $\dim \varphi \leq \tilde{u}(F)$, and if φ_K is isotropic then there exists $\psi \in WF$ with $\dim \psi < \dim \varphi$ such that $\psi_K = \varphi_K$ in WK . This then establishes the excellence of K/F .

Now if $\dim \varphi \leq 3$ or $\varphi \in GPF$ then for any field extension K/F we have that $(\varphi_K)_{\text{an}}$ is defined over F . We thus are done with (i). To establish (ii) (resp. (iii)), it suffices to consider an anisotropic form $\varphi \simeq \langle d, a, b, ab \rangle \in WF$ with φ_K isotropic and show that $(\varphi_K)_{\text{an}}$ is defined over F . If $\varphi_K = 0$ there is nothing else to show, so we may assume $d \notin K^2$. First, consider (ii). Since $\tilde{u}(F) \leq 4$, the $\{\rho_i\} \subset P_2 F$ form a linked group Φ of 2-fold Pfister forms. Now over $E = F(\sqrt{d})$ we have $\varphi_E \in P_2 E$. Clearly, φ becomes isotropic and hence hyperbolic over $E(\{\rho_i\})$ and by LGT there exists $\rho \simeq \langle\langle u, v \rangle\rangle \in \Phi$ such that $\varphi_E \simeq \rho_E$. But then also $\varphi_E \simeq \langle d, a, b, ab \rangle_E \simeq \langle d, u, v, uv \rangle_E$ and by [W, Theorem 7] it follows that φ and $\langle d, a, b, ab \rangle$ are similar over F . In particular, φ contains a subform similar to $\langle u, v, uv \rangle$ which is a Pfister neighbor of $\rho \in \Phi \subset W(K/F)$. Thus, for suitable $x \in \dot{F}$ and with $\psi \simeq (\varphi \perp x\rho)_{\text{an}}$ we get $\dim \psi = 2 < \dim \varphi$ and $\psi_K = \varphi_K \perp x\rho_K = \varphi_K$ in WK , and we are done.

To prove (iii), let $\varphi \simeq \langle d, a, b, ab \rangle$ be as above with φ_K isotropic but not hyperbolic, so that again $d \notin K^2$ and φ_K is similar to $\langle 1, -d \rangle_K$. Over $L = K(\langle 1, -d \rangle) = F(\{\rho_i\}, \langle 1, -d \rangle)$ we have $\varphi_L = 0$. We know from Corollary 4.5 that F is amenable. Thus, there are $\tau, \tau_i \in WF$ such that $\varphi = \sum_i \tau_i \otimes \rho_i + \tau \otimes \langle 1, -d \rangle$ in WF . Now over K we have $\varphi_K = \tau_K \otimes \langle 1, -d \rangle_K$ in WK , and as $d \notin K^2$ we must have that $\dim \tau$ is odd. Choose $b \in \dot{F}$ such that $\tau \perp \langle -b \rangle \in I^2 F$. We then get that $(\tau \perp \langle -b \rangle) \otimes \langle 1, -d \rangle \in I^3 F$. Thus,

$$(\varphi \perp -b\langle 1, -d \rangle)_K = (\tau \perp \langle -b \rangle)_K \otimes \langle 1, -d \rangle_K \in I^3 K .$$

But $\dim(\varphi \perp -b\langle 1, -d \rangle) = 6$ and therefore, APH implies that $(\varphi \perp -b\langle 1, -d \rangle)_K = 0$ and thus $\varphi_K = b\langle 1, -d \rangle_K$ in WK . (iii) now follows.

If $\tilde{u}(F) \leq 6$ then F is 3-linked and therefore $\{\rho_i\} \subset P_3 F$ generates a linked group Φ of 3-fold Pfister forms. In this case, it suffices to consider the case $\dim \varphi \leq 6$. By invoking Proposition 4.7(i), we can conclude that there exists $\pi \in W(K/F) \cap GP_3 F$ such that $\psi \simeq (\varphi \perp -\pi)_{\text{an}}$ has dimension ≤ 4 . But then ψ_K stays anisotropic (its dimension is ≤ 4 and K is a function field of forms of dimension 8!). Clearly, $\psi_K = \varphi_K$ and the anisotropic part of φ_K is therefore defined over F by ψ_K . This establishes (iv).

In (v), excellence of K/F follows in a similar manner from Proposition 4.7(iii). \square

Corollary 4.8 *Let F be a field such that $\bar{u}(F) \leq 6$ or F is linked. Then $F(\pi)/F$ is excellent for any Pfister form π over F .*

5 K -minimal forms

Let K/F be a field extension. In [H 1], the notion of a K -minimal form has been introduced. A form φ over F is said to be K -minimal if φ is anisotropic, φ_K is isotropic, and η_K is anisotropic for any subform $\eta \subset \varphi$ with $\dim \eta < \dim \varphi$. In general, the characterization of K -minimal forms seems to be a rather difficult problem so that it is already quite desirable just to have some information on bounds for the dimensions of K -minimal forms. This leads to the definition of the following two invariants t_{\max} and t_{\min} of the field extension K/F which have been introduced in [H 1] :

$$\begin{aligned} t_{\max}(K/F) &= \sup\{\dim \varphi \mid \varphi \in WF \text{ is } K\text{-minimal}\} \\ t_{\min}(K/F) &= \min\{\dim \varphi \mid \varphi \in WF \text{ is } K\text{-minimal}\} \end{aligned}$$

We put $t_{\min}(K/F) = t_{\max}(K/F) = 1$ if there are no K -minimal forms, i.e., if all anisotropic forms over F stay anisotropic over K .

Our interest is focused on the case where K/F is a Pfister extension. As an easy example, let $\rho \simeq \langle 1, -d \rangle \in P_1F$ be anisotropic and let $K = F(\rho) = F(\sqrt{d})$. Then the K -minimal forms over F are exactly the forms similar to ρ . In particular, $t_{\min}(K/F) = t_{\max}(K/F) = 2$ (see [L 1, Ch.VII, 3.1], or [S, Ch.2, 5.1]). The situation is considerably more complicated in the case where $K = F(\rho)$ with an anisotropic $\rho \in P_2F$. A characterization of K -minimal forms in this case can be found in [HLVG], and in [HVG] it has been dealt with the question how properties of F might affect $t_{\max}(K/F)$. In particular, an example of a field F and an anisotropic $\rho \in P_2F$ has been constructed such that $t_{\max}(K/F) = \infty$.

Generally, very little is known regarding K -minimal forms when $K = F(\rho)$ for an anisotropic $\rho \in P_nF$ if $n \geq 3$. In these cases, we do not even have that K/F is excellent in general. In [I], beside these non-excellence results one can also find examples of such F and ρ where there exist 2^n -dimensional K -minimal forms. These examples have been generalized in [H 6].

We do have the following.

Theorem 5.1 *Let $\emptyset \neq \{\rho_i\} \subset P_nF$ be a set of anisotropic Pfister forms and let $L = F(\{\rho_i\})$. Let K/L with $t_{\min}(K/L) = 1$ or $t_{\min}(K/L) > 2^{n-1} + 1$. Then $t_{\min}(K/F) = 2^{n-1} + 1$. Furthermore, if the set $\{\rho_i\}$ generates a linked group Φ of n -fold Pfister forms then an anisotropic form $\varphi \in WF$ with $\dim \varphi = 2^{n-1} + 1$ is K -minimal iff φ is a Pfister neighbor of some $\rho \in \Phi$. In particular, if $\{\rho_i\} = \{\rho\}$ is a singleton set then the K -minimal forms of dimension $2^{n-1} + 1$ are exactly the Pfister neighbors of ρ of dimension $2^{n-1} + 1$.*

Proof. Any anisotropic form φ over F of dimension $\leq 2^{n-1} + 1$ which becomes isotropic over K must already become isotropic over L . By the Anisotropy Criterion we have $t_{\min}(L/F) \geq 2^{n-1} + 1$ and thus $t_{\min}(K/F) \geq 2^{n-1} + 1$. However, clearly any Pfister neighbor of dimension $2^{n-1} + 1$ of any Pfister form in $\{\rho_i\}$ becomes isotropic over K . Hence, $t_{\min}(K/F) = 2^{n-1} + 1$.

To show the second part of the theorem, it suffices to show that if $\varphi \in WF$ is anisotropic and $\dim \varphi = 2^{n-1} + 1$, and if φ_K is isotropic, then φ is a Pfister neighbor of some $\rho \in \Phi$. So let φ be as above. By the above, φ already becomes isotropic over $L = F(\{\rho_i\})$. In [H 4, Theorem 3], it was shown that if $\{\rho_i\} = \{\rho\}$ is a singleton set then φ_L being isotropic implies that φ is a Pfister neighbor of ρ . The proof of this statement given in [H 4] readily

carries over to the general case using the assumption that the set $\{\rho_i\}$ generates a linked group of Pfister forms and then invoking LGT. This then leads to the desired conclusion that there exists $\rho \in \Phi$ such that φ is a Pfister neighbor of ρ . We omit the details. \square

This theorem applies in particular to the situation where $K = F(\{\rho_i\}, \{\pi_j\})$ with $\{\pi_j\} \subset PF$ a set of anisotropic Pfister forms of fold $> n$. In this case, $K = L(\{\pi_j\})$ and the Anisotropy Criterion implies that either $t_{\min}(K/L) = 1$ or $t_{\min}(K/L) > 2^n$.

Remark 5.2 Suppose the ρ_i 's in the above theorem generate a linked group of Pfister forms. If $n = 2$, one can show that there are no 4-dimensional minimal forms (this is implicit in our proof of Theorem 4.6(ii)), and if $n = 3$, it follows from Proposition 4.7 that there are no 6-dimensional minimal forms. But as mentioned above, there can be 8-dimensional minimal forms.

The next theorem provides upper bounds for a Pfister extension K/F under the additional assumption that $\tilde{u}(F) < \infty$.

Theorem 5.3 *Let $k \geq 1$ and let $\{\pi_\alpha\} \subset PF$ be a set of anisotropic Pfister forms of dimension $\leq 2^k$ such that $\{\pi_\alpha\} \cap P_k F \neq \emptyset$. Let $K = F(\{\pi_\alpha\})$. Then $t_{\max}(K/F) \leq \max\{\tilde{u}(F), 2^k + 1\}$.*

Furthermore, if $\{\pi_\alpha\} \cap P_k F = \{\rho\}$ is a singleton set then $t_{\max}(K/F) \leq \max\{\tilde{u}(F), 2^{k-1} + 1\}$.

Proof. We clearly may assume that $\tilde{u}(F) < \infty$, say $\tilde{u}(F) \leq 2^\ell$ for some ℓ . We also may assume that F is formally real because otherwise there are no anisotropic forms over F of dimension $> \tilde{u}(F)$.

Let $\varphi \in WF$ be anisotropic with $\dim \varphi > \tilde{u}(F)$ and φ_K isotropic. Then by Lemma 3.2, φ contains a Pfister neighbor γ of dimension $2^{n-1} + 1$ of some $\pi \in P_n F \cap W(K/F)$ for some $n \geq 1$. Obviously, γ_K is isotropic because $\pi_K = 0$. To show the first part of the theorem, it therefore suffices to show that if $n - 1 > \max\{k, \ell - 1\}$ then γ is not K -minimal.

So let $n - 1 > \max\{k, \ell - 1\}$, let $\pi \in P_n F \cap W(K/F)$ be anisotropic and let γ be a Pfister neighbor of π . As $\tilde{u}(F) < \infty$ we have that F is SAP, so we may assume that after scaling we have $\text{sgn}_P \gamma \geq 1$ for all $P \in X_F$ (note that $\dim \gamma$ is odd because $n - 1 \geq 1$). By Lemmas 3.1 and 3.2, there exist an integer $r > 0$, $t_i \in \dot{F}$, $\rho_i \in \{\pi_\alpha\}$, $1 \leq i \leq r$, such that for $\tau_i \simeq \langle\langle 1, \dots, 1, t_i \rangle\rangle \in P_{n-m(i)} F$ (where $m(i) \leq k$ denotes the fold of ρ_i), we have that

$$\pi \equiv \sum_{i=1}^r \tau_i \otimes \rho_i \pmod{I^{n+1} F}.$$

Note that $n - m(i) \geq n - k \geq 2$. Thus, let $\tilde{\tau}_i \simeq \langle\langle 1, \dots, 1, t_i \rangle\rangle \in P_{n-m(i)-1} F$ so that $\tau_i \simeq \langle\langle 1 \rangle\rangle \otimes \tilde{\tau}_i$. As $\tilde{u}(F) \leq 2^\ell \leq 2^{n-1}$, we have that F is $(n-1)$ -linked. Hence, there exists $\tilde{\pi} \in P_{n-1} F$ such that

$$\tilde{\pi} \equiv \sum_{i=1}^r \tilde{\tau}_i \otimes \rho_i \pmod{I^n F}.$$

We observe that $\tilde{\pi}_K \equiv 0 \pmod{I^n K}$. Thus, by APH, $\tilde{\pi}_K = 0$, i.e., $\tilde{\pi} \in P_{n-1} F \cap W(K/F)$. Now $\text{sgn}_P(\tau_i \otimes \rho_i) = \text{sgn}_P(\langle\langle 1 \rangle\rangle \otimes \tilde{\tau}_i \otimes \rho_i) = 2\text{sgn}_P(\tilde{\tau}_i \otimes \rho_i)$ for all $P \in X_F$, and one easily checks that therefore

$$\text{sgn}_P \pi = 2\text{sgn}_P \tilde{\pi} \text{ for all } P \in X_F.$$

In particular, γ is indefinite at P (i.e., $1 \leq \text{sgn}_P \gamma \leq 2^{n-1} - 1$) iff $\text{sgn}_P \pi = 0$ iff $\text{sgn}_P \tilde{\pi} = 0$, and γ is definite at P (i.e., $\text{sgn}_P \gamma = 2^{n-1} + 1$) iff $\text{sgn}_P \pi = 2^n$ iff $\text{sgn}_P \tilde{\pi} = 2^{n-1}$. It follows immediately that

$$|\text{sgn}_P(\gamma \perp -\tilde{\pi})| \leq 2^{n-1} - 1 \text{ for all } P \in X_F,$$

and since $\tilde{u}(F) \leq 2^{n-1}$ it follows from Lemma 2.1 that $\dim(\gamma \perp -\tilde{\pi})_{\text{an}} \leq 2^{n-1} - 1$ (recall that $n \geq 2$ and thus $\dim(\gamma \perp -\tilde{\pi})$ is odd). Comparing this with the dimension of $\gamma \perp -\tilde{\pi}$ itself, we get for the Witt index $i_W(\gamma \perp -\tilde{\pi}) \geq 2^{n-2} + 1$. It follows that γ contains a Pfister neighbor $\tilde{\gamma}$ of $\tilde{\pi}$ with $\dim \tilde{\gamma} = 2^{n-2} + 1 < \dim \gamma$. $\tilde{\gamma}_K$ is isotropic because $\tilde{\pi}_K = 0$ and we conclude that γ is not K -minimal.

To prove the remaining part of the theorem, let now $\{\pi_\alpha\} \cap P_k F = \{\rho\}$. Let $\pi \in P_n F \cap W(K/F)$ and let γ be a Pfister neighbor of π with $\dim \gamma = 2^{n-1} + 1$. Again, we assume that $\text{sgn}_P \gamma \geq 1$ for all $P \in X_F$. By a similar reasoning as above, in order to establish that $t_{\max}(K/F) \leq \max\{\tilde{u}(F), 2^{k-1} + 1\}$, it suffices to show that if $n - 1 > \max\{k - 1, \ell - 1\}$ then γ is not K -minimal (as before, ℓ is such that $\tilde{u}(F) \leq 2^\ell$).

So let us assume that $n - 1 > \max\{k - 1, \ell - 1\}$. In fact, if $n - 1 > k$ then the above proof of the first part of the theorem carries over without any changes. Hence, we assume from now on that $n - 1 = k$ and thus necessarily $k \geq \ell$. Again, let us write

$$\pi \equiv \sum_{i=1}^r \tau_i \otimes \rho_i \pmod{I^{n+1}F}$$

with τ_i, ρ_i , and also t_i and $m(i)$ as above. If $m(i) \leq k - 1$ for all $1 \leq i \leq r$ then the proof given above for the first part of the theorem also applies in this situation and we are done. So we may assume that at least one of the ρ_i 's has fold k . By the proof of Lemma 3.1, we may also assume that the ρ_i 's are pairwise non-isometric. Since there is only one k -fold Pfister form ρ in $\{\pi_\alpha\}$, we may therefore assume that $\rho_r \simeq \rho$, $m(r) = k$, and $m(i) \leq k - 1$ for $1 \leq i \leq r - 1$. We put $t_r = t$ and get that $\tau_r \simeq \langle\langle t \rangle\rangle$ (recall that $n - 1 = k$ and thus $n - m(r) = n - k = 1$). We now define the following clopen sets in X_F :

$$\begin{aligned} Y &= \{P \in X_F \mid \pi \text{ is definite at } P\} \\ Y_1 &= \{P \in Y \mid \rho \text{ is definite at } P \text{ and } t <_P 0\} \\ Y_2 &= \{P \in Y \mid \rho \text{ is definite at } P \text{ and } t >_P 0\} \\ Y_3 &= \{P \in Y \mid \rho \text{ is indefinite at } P\}. \end{aligned}$$

Comparing signatures, one easily checks that

$$\text{sgn}_P \langle\langle t \rangle\rangle \otimes \rho = \begin{cases} 0 & \text{if } P \in Y_1 \cup Y_3 \\ 2^n & \text{if } P \in Y_2 \end{cases}$$

and

$$\text{sgn}_P \sum_{i=1}^{r-1} \tau_i \otimes \rho_i \equiv \begin{cases} 2^n \pmod{2^{n+1}} & \text{if } P \in Y_1 \cup Y_3 \\ 0 \pmod{2^{n+1}} & \text{if } P \in Y_2. \end{cases}$$

Let $Z_i = \{P \in Y_1 \mid t_i >_P 0\}$ for $1 \leq i \leq r - 1$ and let $x_i \in \dot{F}$ such that $x_i <_P 0$ if $P \in Z_i$ and $x_i >_P 0$ if $P \notin Z_i$. Finally, let $\tilde{\tau}_i \simeq \langle\langle 1, \dots, x_i t_i \rangle\rangle \in P_{n-m(i)-1} F$ for $1 \leq i \leq r - 1$. One readily concludes that for $1 \leq i \leq r - 1$ we get

$$\text{sgn}_P \tilde{\tau}_i \otimes \rho_i = \begin{cases} 0 & \text{if } P \in Y_1 \\ \frac{1}{2} \text{sgn}_P \tau_i \otimes \rho_i & \text{if } P \notin Y_1 \end{cases}$$

and furthermore

$$\operatorname{sgn}_P \rho = \begin{cases} 2^{n-1} & \text{if } P \in Y_1 \\ \frac{1}{2} \operatorname{sgn}_P \langle\langle t \rangle\rangle \otimes \rho & \text{if } P \in Y \setminus Y_1. \end{cases}$$

As F is $(n-1)$ -linked, there exists $\tilde{\pi} \in P_{n-1}F$ with

$$\tilde{\pi} \equiv \sum_{i=1}^{r-1} \tilde{\tau}_i \otimes \rho_i + \rho \pmod{I^n F}.$$

We note that $\tilde{\pi} \in P_{n-1}F \cap W(K/F)$. Comparing signatures modulo 2^n , one readily concludes that for all $P \in Y$ we have $\operatorname{sgn}_P \tilde{\pi} = \frac{1}{2} \operatorname{sgn}_P \pi = 2^{n-1} = 2^k$. Similarly as above, we see that $\operatorname{sgn}_P \gamma = 2^{n-1} + 1$ iff $\operatorname{sgn}_P \pi = 2^n$, i.e., π is definite at P , which by definition means that $P \in Y$ and thus $\operatorname{sgn}_P \tilde{\pi} = 2^{n-1}$. As $\operatorname{sgn}_P \tilde{\pi} \in \{0, 2^{n-1}\}$ and $1 \leq \operatorname{sgn}_P \gamma \leq 2^{n-1} - 1$ for all $P \notin Y$, we get

$$|\operatorname{sgn}_P(\gamma \perp -\tilde{\pi})| \leq 2^{n-1} - 1 \text{ for all } P \in X_F.$$

Using the same reasoning as before, there exists a Pfister neighbor $\tilde{\gamma}$ of $\tilde{\pi}$ of dimension $2^{n-2} + 1$ with $\tilde{\gamma} \subset \gamma$. $\tilde{\gamma}_K$ is isotropic because $\tilde{\pi}_K = 0$, and $\dim \tilde{\gamma} = 2^{n-2} + 1 < \dim \gamma = 2^{n-1} + 1$, which yields that γ is not K -minimal. \square

Corollary 5.4 *Let F be linked and $\pi \in P_n F$ be anisotropic. Let $K = F(\pi)$. Then the K -minimal forms are exactly the Pfister neighbors of π of dimension $2^{n-1} + 1$.*

Proof. The case $n = 1$ has been mentioned before (and we don't need F being linked in this case). The case $n = 2$ was first proved in [H1, Theorem 4.5.2], a simplified proof can be found in [HLVG, Prop. 4.4]. If $n > 3$ the statement follows from the fact that if F is linked then $\tilde{u}(F) \leq 8$ and by invoking Theorems 5.1 and 5.3. Finally, if $n = 3$ then Theorem 5.3 implies that any K -minimal form has dimension ≤ 8 . However, any anisotropic form over F of dimension ≤ 8 which becomes isotropic over K contains a Pfister neighbor of π of dimension 5 by Proposition 4.7, which completes the proof. \square

6 Characterization of K -minimal forms

In the previous section, we have provided certain bounds for the dimensions of K -minimal forms for Pfister extensions K/F and found an exact lower bound in all cases, and upper bounds under the assumption that $\tilde{u}(F) < \infty$. We now will give a precise description of K -minimal forms for Pfister extensions K/F where $\tilde{u}(F) < \infty$ and where the Pfister forms involved are of "suitably" large dimension.

It follows easily from the definition of K -minimal forms that anisotropic Pfister forms of fold ≥ 2 can never be K -minimal for any field extension K/F . This is simply because if π_K is isotropic then π_K is hyperbolic and any subform of π of dimension $2^{n-1} + 1 < 2^n = \dim \pi$ will be isotropic over K . Nevertheless, we will define a version of K -minimality for Pfister forms which will be useful later on.

Definition 6.1 Let K/F be a field extension. Then $\pi \in P_n F$, $n \geq 1$, is a *Pfister K -minimal form* if the following three conditions are fulfilled :

- (i) π is anisotropic over F .
- (ii) π_K is isotropic and hence hyperbolic, i.e., $\pi \in W(K/F)$.
- (iii) For any $\sigma \in P_m F$ with $\sigma \subset \pi$ and $m < n$ we have that σ_K is anisotropic.

One easily sees that to check whether condition (iii) holds it suffices to check that it holds for $m = n - 1$. In this case ($m = n - 1$) and with σ and π as above, there exists $t \in \dot{F}$ such that $\pi \simeq \sigma \otimes \langle\langle t \rangle\rangle$.

We would like to describe the K -minimal forms for some of the Pfister extensions K/F dealt with in the previous sections. It turns out that they will be Pfister neighbors. Indeed, they will be Pfister neighbors of Pfister K -minimal forms. The following lemma relates K -minimal forms which are Pfister neighbors and Pfister K -minimal forms. The assumptions are in a way so that they apply directly to the cases we consider.

Lemma 6.2 *Let F be formally real with $\tilde{u}(F) \leq 2^n$. Let K/F be a field extension such that K -minimal forms are always Pfister neighbors. Let $\pi \in P_{n+1}F \cap W(K/F)$ be anisotropic and let φ be a Pfister neighbor of π . Then the following statements are equivalent.*

- (i) φ is K -minimal.
- (ii) $\dim \varphi = 2^n + 1$ and π is Pfister K -minimal.

Proof. The case $n = 0$ is trivial. So let $n \geq 1$. Also, we may assume from the beginning that $\dim \varphi = 2^n + 1$. For if a Pfister neighbor φ of the $(n + 1)$ -Pfister form π is isotropic, then π is hyperbolic and any subform of φ of dimension $2^n + 1$ is isotropic as well.

(i) \Rightarrow (ii). Suppose π is not Pfister K -minimal. Then there exist $\sigma \in P_n F$, $t \in \dot{F}$, such that $\sigma_K = 0$ and $\pi \simeq \sigma \otimes \langle\langle t \rangle\rangle$. Let $Y = \{P \in X_F \mid \text{sgn}_P \sigma = 0\}$ and $Z = \{P \in X_F \mid \text{sgn}_P \pi = 0\}$. Note that $Y \subset Z$, and that $\text{sgn}_P \sigma = 2^n$ for all $P \in X_F \setminus Y$ and $\text{sgn}_P \pi = 2^{n+1}$ for all $P \in X_F \setminus Z$. As $\tilde{u}(F) < \infty$, F is SAP and we may assume that after possibly scaling, we have $\text{sgn}_P \varphi \geq 1$ for all $P \in X_F$ (note that $\dim \varphi$ is odd). In particular, $\text{sgn}_P \varphi = 2^n + 1$ for all $P \in X_F \setminus Z$ and $1 \leq \text{sgn}_P \varphi \leq 2^n - 1$ for all $P \in Z$. We get

$$|\text{sgn}_P(\varphi \perp -\sigma)| \leq 2^n - 1 \text{ for all } P \in X_F.$$

Hence, by Lemma 2.1 and with $\tilde{u}(F) \leq 2^n$, it follows that $\dim(\varphi \perp -\sigma)_{\text{an}} \leq 2^n - 1$ (note that $\dim(\varphi \perp -\sigma)_{\text{an}}$ is odd). By comparing dimensions, we conclude that $i_W(\varphi \perp -\sigma) \geq 2^{n-1} + 1$. Therefore, there exists $\chi \subset \sigma$ with $\dim \chi = 2^{n-1} + 1$ such that $\chi \subset \varphi$. In particular, χ is a Pfister neighbor of σ , and because $\sigma_K = 0$ we have that χ_K is isotropic. But $\chi \subset \varphi$ and $\dim \chi \leq \dim \sigma < \dim \varphi$ which implies that φ is not K -minimal.

(ii) \Rightarrow (i). As $\pi_K = 0$ we have that φ_K is isotropic. If φ is not K -minimal then there exists a K -minimal form $\chi \subset \varphi$ with $\dim \chi < \dim \varphi = 2^n + 1$. By assumption, χ itself is a Pfister neighbor. Thus, χ is a Pfister neighbor of some $\tau \in P_m F$ with $m \leq n$. Then $\chi_{F(\tau)}$ is isotropic, hence also $\varphi_{F(\tau)}$ and therefore $\pi_{F(\tau)}$ is hyperbolic. We readily conclude that by the Cassels-Pfister subform theorem, $\tau \subset \pi$. Also, χ_K is isotropic and thus τ_K is hyperbolic. In particular, π is not Pfister K -minimal. \square

It should be noted that the assumption on F being formally real with $\tilde{u}(F) \leq 2^n$ was only needed to show (i) \Rightarrow (ii), and that the assumption on K -minimal forms being Pfister neighbors was only used to show (ii) \Rightarrow (i).

Remark 6.3 There are examples of field extensions K/F where there exists a Pfister K -minimal form π which contains K -minimal forms of dimension $\leq \frac{1}{2} \dim \pi$. For instance, let F be any field with an anisotropic 3-fold Pfister form π such that π contains a subform ψ with $\dim \psi = 4$ and $\psi \notin GP_2 F$. Let $K = F(\psi)$. The only anisotropic forms of dimension ≤ 4 over F which become isotropic over K are exactly the forms similar to ψ (see, e.g., [H 2]). It follows that ψ , which is not a Pfister neighbor, is K -minimal, and that π is

Pfister K -minimal. (ii) $\not\Rightarrow$ (i) in this case because no 5-dimensional subform of π containing ψ will be K -minimal. Note that in this example, K/F is not excellent because ψ is not a Pfister neighbor.

Remark 6.4 If all K -minimal forms are Pfister neighbors then one readily sees that K/F is excellent. However, the converse is not true in general. In [HVG], there is an example of a field F with an anisotropic conic $\rho \simeq \langle 1, a, b \rangle$ over F such that for $K = F(\rho)$ one has $t_{\max}(K/F) = \infty$. In particular, by [HLVG, Corollary 3.4], there exist 7-dimensional K -minimal forms. Such a form cannot be a Pfister neighbor because if φ is an anisotropic 7-dimensional Pfister neighbor over F with φ_K isotropic then it is an easy exercise to show that φ contains a subform similar to ρ and is therefore not K -minimal. Hence, there exist (7-dimensional) K -minimal forms which are not Pfister neighbors. On the other hand, K/F is known to be excellent (see the introduction).

For the remainder of this section, we fix the following notations.

m	an integer ≥ 1
F	a field ($\text{char } F \neq 2$) which fulfills the following:
	If $m = 1$ or 2 then $\tilde{u}(F) \leq 2^m$.
	If $m = 3$ then $\tilde{u}(F) \leq 6$ or $\tilde{u}(F) = 8$ and F linked.
	If $m \geq 4$ then $\tilde{u}(F) \leq 2^{m-1}$
$\{\rho_i^{(n)}\}, n \geq m$	a set of n -fold anisotropic Pfister forms over F , possibly empty if $n > m$, but with $\{\rho_i^{(m)}\} \neq \emptyset$.
Φ_m	the linked group of m -fold Pfister forms generated by $\{\rho_i^{(m)}\}$.
Φ'_m	the set $\{2^{m-1}\mathbf{H}\}$ consisting only of the hyperbolic m -fold Pfister form.
$\Phi'_n, n > m$	the set $\{\pi \otimes \langle\langle t \rangle\rangle \mid \pi \in \Phi_{n-1}, t \in F\}$ of n -fold Pfister forms.
$\Phi_n, n > m$	the linked group of n -fold Pfister forms generated by the sets Φ'_n and $\{\rho_i^{(n)}\}$.

Theorem 6.5 *With the notations as above, let $K = F(\{\rho_i^{(n)}\}, n \geq m)$. Let $\varphi \in WF$ be anisotropic. Then the following statements are equivalent.*

- (i) φ is K -minimal.
- (ii) There exist $n \geq m$ and $\pi \in \Phi_n \setminus \Phi'_n$ such that φ is a Pfister neighbor of dimension $2^{n-1} + 1$ of π .

Proof. It follows from Lemma 3.2 and Proposition 4.7 (see also Remark 5.2 in the case $m = 2$), that any anisotropic form over F which becomes isotropic over K already contains a Pfister neighbor which becomes isotropic over K . In particular, K -minimal forms will always be Pfister neighbors. Now if φ is a Pfister neighbor of some n -fold Pfister form π and if φ is K -minimal, then clearly $\dim \varphi = 2^{n-1} + 1$ and necessarily $n \geq m$ by the Anisotropy Criterion. Thus, K -minimal forms are Pfister neighbors of dimension $2^{n-1} + 1$ where $n \geq m$.

The K -minimal forms of dimension $2^{m-1} + 1$ are exactly the Pfister neighbors of dimension $2^{m-1} + 1$ of anisotropic Pfister forms in Φ_m (i.e., Pfister forms in $\Phi_m \setminus \Phi'_m$) by Theorem 5.1.

